



- Figuring the shellfish farming sector

The industry is considered as a whole (at a first stage), but several exploitations systems may be also considered (various specialisations)

Typology of exploitation systems

- Rationale of the approach

- . Oyster production (at each stage: spat collection, growing) depends on natural environment factors (drivers), esp. river flows (salinity, nutrients)
 - ⇒ Link with the coastal water productivity module
- . Producers target an \ll objective function \gg = maintaining the current mean annual production over time (recent observations support this assumption)
- . Producers decisions: if the current production differs from the objectives, the producers may import (export) the equivalent of the negative (positive) differential in order to minimize (maximize) commercial losses (gains)


- Expected output of the shellfish farming module

The model will estimate the gaps between real production and objectives:

- . frequency and intensity of required adaptations
- . additional costs (benefits) associated with the productivity losses (gains) due to freshwater relative scarcity (abundance)

□⇒ Damage costs assessment







Grow-out start:

- size
- population



Oyster Growth DEB Model:

- Temperature
- Food (Phytoplankton)



Natural mortality




Harvest of marketable sized oysters:
Production target



Oyster growth

Sub cohort variability: Harvest Xk DEB model factor season


Length (cm)





Oyster standing stock






Time (days)



Output: Sales and targets (in tons)

Harvest to be sold






Time (days)



Adaptative management





Shellfish Farm specialization

Lately, shellfish farmers have been specializing their activity. The model will represent several types of oyster exploitation.

The shellfish farm block has been developped as a generic block.





Cédric Bacher, Johanna Béganton, Rémi Mongruel, José A. Pérez Agúndez, Jean Prou

